

ΤΟ ΒΗΜΑ ΤΟΥ ΑΣΚΛΗΠΙΟΥ

VEMA OF ASKLIPIOS

APRIL-JUNE 2002 VOLUME 1 No 2


QUARTERLY EDITION BY THE 1st NURSING DEPARTMENT
OF ATHENS TECHNOLOGICAL EDUCATIONAL INSTITUTION

Ιστορική εξέλιξη του φαρμάκου
Φροντίδα ηλικιωμένων
Υποδοχή ασθενών στο Τμήμα Επειγόντων Περιστατικών
Συχνότητα κατακλίσεων σε επαρχιακό νοσοκομείο
Β-θαλασσαιμία και φυσιολογικός καταμήνιος κύκλος
Εντερική θρέψη σε εγκαυματίες και πολυτραυματίες
Νοσηλευτική θεωρία της πολυπλοκότητας
Αξιοποίηση του χρόνου εργασίας

Historical evolution of medicines
Elderly care
Patient's reception in Emergency Department
The frequency of pressure sores in a provincial hospital
B-thalassaemia and the frequency of the menstrual cycle
Enteral nutrition in burnt and multiple trauma patients
Nursing theory of the complexity
Effective time management in the working place



ΕΛΛΑΣ

ΕΛΛΑΣ
ΑΘΗΝΑ
2671
ΕΠΙΧΕΙΡΗΣΗHELLAS
PAYE
PORT

ΕΝΤΥΠΟ ΚΛΕΙΣΤΟ ΑΡ. ΑΔΕΙΑΣ 1459/99

ΒΗΜΑ ΙΑΤΡΙΚΗΣ ΕΚΔΟΣΕΙΣ
Κατεχάκη & Αφριανείου 3 - 115 25 ΑΘΗΝΑ

Περιεχόμενα

Ανασκοπήσεις

Ιστορική εξέλιξη του φαρμάκου. Γ.Α. Φούκα 57

Φροντίδα ηλικιωμένων και επαγγελματική εξουθένωση. Ε. Κοτρώτσιου, Θ. Παραλίκας 61

Ερευνητικές εργασίες

Υποδοχή ασθενών στο Τμήμα Επειγόντων Περιστατικών της Ευρωκλινικής Αθηνών. Ν.Β. Φώτος, Π. Επιτροπάκης 65

Μέτρηση συχνότητας και βαρύτητας κατακλίσεων σε ασθενείς ελληνικού επαρχιακού γενικού νοσοκομείου. Ι. Παπαθανασίου, Θ. Παραλίκας, Ε. Λαχανά, Γ. Τζαβέλας, Σ. Κοτρώτσιου, Β. Κουτσοπούλου 72

Β-θαλασσαιμία και φυσιολογικός καταμήνιος κύκλος. Μπορούμε να αναμένουμε ένα καλύτερο κύκλο αν τα επίπεδα φερριτίνης διατηρούνται σε ένα χαμηλότερο επίπεδο τιμών; Μ. Γουρνή, Δ. Σαπουντζή-Κρέπια, Ζ. Ρούπα-Δαριβάκη, Μ. Σκάντζος, Ν. Σκορδής, Χ. Μπαρτσόκας, Ε. Θεοδοσοπούλου 77

Ο ρόλος της έγκαιρης εντερικής θρέψης σε εγκαυματίες και πολυτραυματίες. Ομοιότητες και διαφορές. Χρ. Μαρβάκη, Θ. Οικονόμου, Π. Ιορδάνου, Ε. Μαρβάκη, Ε. Θεοδοσοπούλου, Ε. Πατηράκη 83

Ειδικά άρθρα

Μετα-ολιστικό παράδειγμα: Μία νέα αλλαγή παραδείγματος μέσα από την ανάπτυξη της ενοποιημένης νοσηλευτικής θεωρίας της πολυπλοκότητας. S.L. Van Sell, I. Καλοφισούδης 89

Μπορούμε να κερδίσουμε χρόνο βελτιώνοντας τις σχέσεις μας στο χώρο εργασίας; Κ. Τριανταφυλλιάκης 93

Οδηγίες για τους συγγραφείς 99

Contents

Reviews

Historical evolution of medicines. G.A. Fouka 57

Elderly care and the burnout syndrome. E. Kotrotsiou, Th. Paralikas 61

Original papers

Patients' reception in Emergency Department. N.V. Fotos, P. Epitropakis 65

Measurement of the frequency of pressure sores in patients of a Greek general provincial hospital. I. Papathanasiou, Th. Paralikas, E. Lahana, G. Tzavelas, S. Kotrotsiou, V. Kutsopoulou 72

B-thalassaemia and normal menstrual cycle. Would we expect a better menstrual cycle if ferritin levels can be maintained at a lower level? M. Gourni, D. Sapountzi-Krepia, Z. Roupa-Darivaki, M. Sgantzos, N. Skordis, C. Bartsokas, H. Theodossopoulou 77

The role of early enteral nutrition in burnt and multiple trauma patients. Similarities and differences. Ch. Marvaki, T. Iconomou, P. Iordanou, E. Marvaki, E. Theodossopoulou, E. Patiraki 83

Special articles

Metaholistic paradigm. A new paradigm shift through the development of the complexity integration nursing theory. S.L. Van Sell, I. Kalofissudis 89

Can effective time management improve relations in the working place? C. Triadafillakis 93

Instructions to authors 99

The role of early enteral nutrition in burnt and multiple trauma patients

Similarities and differences

Ch. Marvaki,¹
T. Iconomou,²
P. Iordanou,¹
E. Marvaki,³
E. Theodossopoulou,⁴
E. Patiraki⁴

¹Nursing Department of TEI of Athens

²MD, Department of Plastic Surgery, Microsurgery and Burn Center, "G. Gennimatas" General State Hospital, Athens

³MD, Moliere Hospital of Brussels

⁴School of Nursing, University of Athens, Athens, Greece

Aim The aim of this study was to investigate the role of Early Enteral Nutrition (EEN) in burnt and multiple trauma patients. **Material-Method** 31 patients with burns >25% of the total body surface area and 35 patients with more than two traumas were included in our study. A nutritional support protocol was followed. Blood markers and the general condition of patients were assessed for the role of EEN. **Results** In burnt patients, total proteins, serum albumin and serum globulin were increased significantly between all measurements. In multiple trauma patients total proteins and serum globulin were increased significantly between the 1st and 3rd measurement ($P<0.001$). Serum albumin was slightly decreased between the 1st and 3rd measurement ($P=0.350$). For burnt patients, between the 1st and 3rd measurement, Fe was increased ($P<0.008$) and for multiple trauma it was slightly increased. For burnt patients TIBC was increased ($P<0.001$) and for multiple trauma patients was slightly decreased. There were no significant changes in the absolute number of lymphocytes neither in immunoglobulin-IgA, IgM, IgG for both patients groups. The mean serum albumin value of the burnt patient was significantly increased compared with the multiple trauma patients ($P=0.012$). The mean serum globulin value of the multiple trauma patients was significantly increased compared with the burnt patients ($P=0.019$). TIBC did not increase between groups in the 1st and 2nd measurements, but it was significantly increased in the 3rd measurement in burnt patients compared with the multiple trauma patients ($P=0.005$). The mean value of immunoglobulin IgG showed a significant increase in multiple trauma patients compared with the burnt patients ($P=0.006$). **Conclusions** Generally, the results of our study confirm the effectiveness of EEN in both groups of patients as well as the existence of similarities and differences.

Περίληψη Ο ρόλος της έγκαιρης εντερικής θρέψης σε εγκαυματίες και πολυτραυματίες. **Ομοιότητες και διαφορές.** Χρ. Μαρβάκη,¹ Θ. Οικονόμου,² Π. Ιορδάνου,¹ Ε. Μαρβάκη,³ Ε. Θεοδοσοπούλου,⁴ Ε. Πατριάκη.⁴ ¹Τμήμα Νοσηλευτικής TEI Αθénas, ²Τμήμα Πλαστικής Χειρουργικής, Μικροχειρουργικής και Εγκαυμάτων, Νοσοκομείο «Γ. Γεννηματάς», Αθήνα, ³Νοσοκομείο "Moliere", Βρυξέλλες, ⁴Τμήμα Νοσηλευτικής, Πανεπιστήμιο Αθηνών, Αθήνα. Το Βήμα του Ασκληπιού 2002, 1(2):83-88. Σκοπός Σκοπός της μελέτης ήταν να εξετάσει το ρόλο της Έγκαιρης Εντερικής Θρέψης (ΕΕΘ) σε εγκαυματίες και πολυτραυματίες. Υλικό-Μέθοδος Στη μελέτη μας συμπεριλήφθησαν 31 εγκαυματίες με εγκαύματα >25% της ολικής επιφανείας σώματος και 35 πολυτραυματίες με περισσότερα από δύο τραύματα. Ακολούθησε ειδικό πρωτόκολλο θρέψης και εκτιμήθηκαν εργαστηριακές παράμετροι και η γενική κατάσταση των ασθενών. Αποτελέσματα Στους εγκαυματίες, τα οδικά λευκώματα, οι λευκωματίνες και οι σφαιρίνες αυξήθηκαν σημαντικά μεταξύ όλων των μετρήσεων. Στους πολυτραυματίες τα οδικά λευκώματα και οι σφαιρίνες αυξήθηκαν σημαντικά μεταξύ 1ης και 3ης μέτρησης ($P<0,001$). Οι λευκωματίνες μειώθηκαν ελαφρά μεταξύ 1ης και 3ης μέτρησης ($P=0,350$). Στους εγκαυματίες, μεταξύ 1ης και 3ης μέτρησης, ο Fe αυξήθηκε ($P<0,008$). Στους πολυτραυματίες ήταν ελαφρά αυξημένος. Στους εγκαυματίες η δεσμευτική ικανότητα σιδήρου (TIBC) αυξήθηκε ($P<0,001$) και στους πολυτραυματίες μειώθηκε ελαφρά. Δεν υπήρξαν σημαντικές διαφοροποιήσεις στον απόλυτο αριθμό των λεμφοκυττάρων ούτε σε αυτό των ανοσοσφαιρινών IgA, IgM, IgG και για τις δύο ομάδες ασθενών. Μεταξύ των ομάδων, η μέση τιμή των λευκωματινών στους εγκαυματίες αυξήθηκε σημαντικά έναντι των πολυτραυματιών ($P=0,012$). Η μέση τιμή των σφαιρινών στους πολυτραυματίες αυξήθηκε σημαντικά έναντι των εγκαυματιών ($P=0,019$). Η TIBC των δύο ομάδων δεν αυξήθηκε μεταξύ 1ης και 2ης μέτρησης, αλλά αυξήθηκε σημαντικά στην 3η μέτρηση στους εγκαυματίες σε σχέση με τους πολυτραυματίες ($P=0,005$). Η μέση τιμή των ανοσοσφαιρινών IgG παρουσίασε μία σημαντική αύξηση στους πολυτραυματίες συγκριτικά με τους εγκαυματίες ($P=0,006$). Συμπεράσματα Γενικά, τα αποτελέσματα της παρούσας μελέτης επιβεβαίωσαν την αποτελεσματικότητα της ΕΕΘ και στις δύο ομάδες των ασθενών καθώς επίσης και παρουσία ομοιοτήτων και διαφορών μεταξύ των ομάδων.

Key words: Role of early enteral nutrition, nutritional markers (total proteins, blood albumin, globulin, absolute number of lymphocytes, immunoglobulin-IgA, IgM, IgG, Fe (iron) and TIBC-Total Iron Binding Capacity)

Λέξεις κλειδιά: Ρόλος έγκαιρης εντερικής θρέψης, εργαστηριακές παράμετροι θρέψης (οδικά λευκώματα, λευκωματίνη ορού, σφαιρίνες, απόλυτος αριθμός λεμφοκυττάρων, ανοσοσφαιρίνες IgA, IgM, IgG, σίδηρος και σιδηροδεσμευτική ικανότητα)

Severely burnt patients generally show a marked hypermetabolic response and their energy expenditure increases to almost twice the normal one as burn size exceeds 50% of TBSA.¹⁻³ Referring to multiple trauma patients, hypermetabolism occurs in most of them within 48 hours after injury.⁴⁻⁶ This hypermetabolic response, for both burnt and multiple trauma patients, is accompanied by a progressive decline of host defences, immunological abnormalities and a marked decline in the circulating number of T lymphocytes that impair survival.⁷⁻¹⁰ Aggressive nutritional support to meet the increased energy expenditure has been considered essential for the management of burnt patients.⁸ Enteral nutrition has been considered as an essential part of trauma management to minimize the catabolic loss and enhance the immunological response.^{11,12} It has been shown, for both burnt and multiple trauma patients, to be an effective additional measure in stress, ulcer prophylaxis and prevention of sepsis.¹³⁻¹⁹ The purpose of our study was firstly to evaluate the effectiveness of EEN in burnt and multiple trauma patients, in association with their post burnt and post trauma nutritional state, their immunologic response and secondly to examine similarities between the two patients' groups.

Material and method

Our study included 31 patients with burns >25% of the total body surface area, who were admitted in the department of plastic surgery and burn unit of the general state hospital (GSH) of Athens, Greece, "G. Gennimatas" and 35 patients with more than two traumas, who were admitted in the Intensive Care Unit of the GSH of Athens, Greece, "KAT" over the last 3 years.

Inclusion criteria for burnt patients were: (a) the age (≥ 17 years) and (b) the extent of the burn injury. Exclusion criteria included the following: (a) patients in palliative care, (b) previous or planned surgical operation, (c) patients with sepsis, (d) patients presenting allergies to the enteral fluids. Inclusion criteria for multiple trauma patients were: (a) the age (≥ 15 years) and (b) a trauma at least in two different organs or organ systems in different parts of the body, with at least one life-threatening organ injury. Exclusion criteria included the following: (a) patients in palliative care, (b) previous or planned surgical operation, (c) patients with sepsis, (d) presenting allergies to the enteral fluids, (e) patients with burn injury, (f) pre-existing diabetes mellitus or renal or hepatic disease and (g) pregnant women. Data were collected using an assessment chart. The two hospital's Ethic Committee approved of the study and all the patients selected gave their consent. A nutritional support protocol was followed in all patients included EEN. It started the first six hours of admission for burnt patients and the first two days of admission for multiple trauma patients (three of the patients started the 5th day). This lasted until complete healing of burn injuries and until complete treatment of multiple trauma patients. Solu-

tions of different nutritional value were administered via nasogastric feeding tube according to the energy requirements and gastrointestinal tolerance of the patients. The enteral nutrition formulas used were: (a) Low kcal 0.5 kcal/mL with N 2 g%. (b) Standard value 1 kcal/mL with N 4 g% and (c) high kcal value 1.5 kcal/mL with N 6 g%.

Results

31 burnt patients mean aged 43.7 years-old (SD 12.2) and 35 multiple trauma patients mean aged 40.2 years-old (SD=18.6) were studied. 18 (58.1%) burnt patients were men and 13 (41.9%) women. 25 (71.4%) multiple trauma patients were men and 10 (28.6%) were women. Characteristics of burnt patients are presented in (tabl. 1). The results in burnt patients were as following: all proteins showed a significant increase between the three measurements such as: Total proteins in the 1st measurement ($P=0.003$), at the 2nd ($P=0.004$) and at the 3rd ($P<0.001$). In multiple trauma patients: total proteins at the 1st measurement ($P=0.024$), at the 2nd ($P=0.010$) and at the 3rd ($P<0.001$) (fig. 1). In burnt patients: serum albumin showed a significant increase at the 1st measurement ($P=0.015$), the 2nd ($P=0.120$) and the 3rd ($P<0.005$). In multiple trauma patients: serum albumin was slightly decreased at the 1st measurement ($P=0.790$), the 2nd mean value was decreased ($P=0.270$) and the 3rd mean value was decreased ($P=0.350$) (fig. 2). For burnt patients: serum globulin was increased significantly at all measurements, at the first measurement ($P=0.05$), at the 2nd ($P=0.007$) and at the 3rd ($P<0.001$). For

Table 1. Characteristics of the 31 burnt patients.

Characteristics	Mean value	SD
Body weight	73.56	10.43
Total body surface area burned %	47.40	18.67




Figure 1. Mean value of total proteins (mg/dL).




Figure 2. Mean value of serum albumin (mg/dL).

multiple trauma patients: serum globulin at the 1st measurement was increased ($P < 0.001$), at the 2nd was slightly increased ($P = 0.570$), and at the 3rd was increased ($P < 0.01$) (fig. 3).

For burnt patients: Fe level did not change significantly at the 1st measurement ($P = 0.09$) neither at the 2nd ($P = 0.280$) but was significantly increased at the 3rd ($P < 0.008$). TIBC was increased significantly at the 1st measurement ($P < 0.008$). TIBC was increased significantly at the 1st measurement ($P < 0.001$) (tabl. 2). There were no significant changes in the absolute number of lymphocytes. The mean value of immunoglobulin IgG was increased in all measurements. The mean value of immunoglobulin IgA was slightly decreased in all measurements. The mean value of immunoglobulin IgM was increased at the 1st and 2nd measurements but was decreased at the 3rd measurement. The mean value and SD of the numbers of lymphocytes, IgG, IgA, IgM, Fe and TIBC are presented in (tabl. 2). Complications related to EEN were diarrhea in 6.8% of our patients hyperglycemia in 19.4% and electrolyte disturbances in 45.2%.




Figure 3. Mean value of serum globulin (mg/dL).

For multiple trauma patients: Fe level did not change significantly between measurements, but was slightly increased (tabl. 2). TIBC almost remained stable (tabl. 2). There were no significant changes in the absolute number of lymphocytes. The mean values of immunoglobulin IgA and IgM were slightly increased in all measurements. The mean value of immunoglobulin IgG was increased in all measurements; at the 1st measurement was increased significantly ($P = 0.004$), at the 2nd was increased ($P = 0.108$), and at the 3rd was significantly increased ($P = 0.017$). The mean value and SD of the numbers of lymphocytes, IgG, IgA, IgM, Fe and TIBC are presented in (tabl. 2). Complications related to EEN were diarrhea in 11.4% of our patients, hyperglycemia in 17.1% and electrolyte disturbances in 34.3%.

Regarding the results between groups patients. There were no significant differences in mean value of total proteins between groups.

The mean serum albumin value of the burnt patients was significantly increased compared with the multiple trauma patients ($P = 0.012$).

The mean serum globulin value of the multiple trauma patients was significantly increased compared with the burnt patients ($P = 0.019$).

Fe level did not change significantly between groups.

TIBC was not increased between groups at the 1st and the 2nd measurements, but it was significantly increased at 3rd measurement in burnt patients compared with the multiple trauma patients ($P = 0.005$). There were no significant changes in the absolute number of lymphocytes between groups in all measurements.

The mean values of immunoglobulin IgA and IgM were higher in multiple trauma patients but there was no significant difference. The mean value of immunoglobulin IgG showed a significant increase in multiple trauma patients compared with the burnt patients ($P = 0.06$).

Discussion

The concept of the administration of EEN nutrition after injury is relatively new and based upon animal experiments performed more than a decade ago. Based on the animal studies, it seemed reasonable that early enteral nutrition would help to prevent the development of inflammatory and infectious complications in seriously injured patients. One of the problems was that very early enteral nutrition was difficult to applied in a great number of trauma patients.²⁰ Gianelli et al²¹ performed

Table 2. Mean value and SD between measurements in some of the markers of 31 burnt and 35 multiple trauma patients.

Markers	Measurements	Mean value	SD	Mean value	SD
		Burnt patients		Multiple trauma patients	
Absolute number of lymphocytes (%)	1st day	1731.3	952	1456.5	980
	8th day	1426.8	677.6	1353.1	625.3
	15th day	1526.1	577.3	1420.8	697.5
IgG (mg/dL)	1st day	922.0	346.6	1270.8	526.62
	8th day	955.7	313.8	1396.8	562.47
	15th day	997.1	308.8	1495.45	488.88
IgA (mg/dL)	1st day	223.82	125.10	205.81	107.69
	8th day	211.30	83.40	204.05	82.37
	15th day	96.90	64.30	203.00	102.87
IgM (mg/dL)	1st day	91.90	61.85	166.67	83.87
	8th day	211.30	83.40	204.05	82.37
	15th day	197.80	79.75	222.05	72.93
Fe ($\mu\text{g/L}$)	1st day	34.56	19.87	49.67	21.46
	8th day	39.25	27.65	53.00	12.12
	15th day	18.11	19.66	53.33	9.87
TIBC ($\mu\text{g/L}$)	1st day	84.80	42.50	48.30	0.6
	8th day	97.80	48.60	48.00	0.0
	15th day	120.60	39.14	48.00	0.0

the first well controlled clinical study that showed early enteral nutrition compared with delayed enteral nutrition had a beneficial effect on seriously burnt patients. Nutritional therapy plays a key role in the overall management of the burnt and multiple trauma patient.²²⁻²⁶ EEN has been demonstrated to minimize the catabolic loss, to enhance the immunologic response,^{5,7} to prevent stress ulcer^{13,17,27} and to prolong survival. The presence of food in the gut has been shown to stimulate mucosal proliferation, probably as a result of both humoral and nutritional mechanisms.^{28,29} Improved mucosal integrity has also been demonstrated in burnt patients who had received EEN. These findings suggest that EEN may help to diminish the incidence and severity of bacteria translocation by improving or preventing breach of the mucosal barrier.

Concerning the immunological status of burnt patients it has been shown that total IgG, IgA, IgM-secreting cells and B-cell number decrease in burn injuries. Both, cellular and humoral immune systems have been reported to be affected in thermal injuries leading to

immunosuppression and sepsis.³⁰ Similar results were presented for multiple trauma patients.³¹ The results of our study showed that the administration of EEN in burnt patients can enhance their immunologic response leading to increased values of immunoglobulins IgG and IgM. Similarly, in multiple trauma patients the administration of EEN can enhance their immunologic response leading to increased values of all immunoglobulins. Although the absolute number of lymphocytes did not increase between measurements, the fact that there was nonmarked decline in the circulating number of lymphocytes indicated that EEN played a significant role in the improvement of the immunological response of the sample.

Studies have suggested that early enteral feeding has been a helpful aid to recovery complications were rare and prolonged enteral feedings were occasionally required in seriously injured patients.³²⁻³⁴ Burns, sepsis, injury or surgery lower serum glutamine levels. It has been suggested that the lowered plasma glutamine concentration contributes, at least in part, to immunosuppression.³⁵

Researchers showed that enteral diet supplements with arginine in burnt rats decreases the mRNA expression of inflammatory cytokines in organs and improves the survival rate.³⁶ In addition, they showed that administration of enteral glutamine decreased the incidence of infectious complications in multiple trauma patients.³⁷ Our results in burnt patients showed that total proteins, as well as serum albumin, presented a significant increase between measurements. In multiple trauma patients total proteins showed a significant increase between measurements while serum albumin slightly decreased. This fact, for both burnt and multiple trauma patients, confirmed the positive effect of EEN on the improvement of the nutritional state of the patients.

Serum iron levels present a great number of fluctuations during the day, depending on different factors that affect iron distribution between plasma and storage organs. TIBC values are also affected by iron fluctuations and should be studied together with serum iron levels. Furthermore, low serum iron concentration is related to sepsis in burnt patients. Belmonte et al³⁷ studied iron metabolism in burnt children and concluded that hyposideraemia is a frequent finding in the acute phase of the burnt injury and is accompanied by increased ferritin levels and decreased transferrin concentrations. He suggested that low iron values tend to recover without the use of iron supplementation because of an endogenous block released in the acute phase; he thus indicated that iron therapy should not be recommended in the initial period of stress in burnt patients. However, the results of our study in burnt patients showed that Fe values were slightly increased at 1st and 2nd measurements but the significant increase was at 3rd measurement. Fe value in multiple trauma patients did not change significantly between measurements, but it was slightly increased.

In our study TIBC was significantly increased at the 2nd and at the 3rd measurements in burnt patients while in multiple trauma patients it almost remained stable. These results indicated the effectiveness of EEN on the sample of our study.

We should also refer to the similarities and the differences of the burnt and multiple trauma patients that both represent severe and complicated types of injury the body can sustain. One of the differences is that the enteral nutrition in burnt patients could be administered within the first six hours of admission while in multiple trauma patients within the first two days of admission. This was due to the fact that it takes more time to define the needs and the tolerance of multiple trauma patients. However there were no significant differences in mean value of total proteins between groups. It

means that enteral nutrition has various potential similar advantages for both patients' groups, burns and multiple trauma, including a reduction in infectious events and improvement of the survival rate as it is already mentioned. In addition, it should be stressed that none of the sample presented sepsis. Fe level did not change significantly between groups, there were no significant changes in the absolute number of lymphocytes between groups in all measurements and the mean value of immunoglobulins was IgA and IgM higher in multiple trauma patients but there was no significant difference. The above results can be explained as an evidence of similarities existing in both patients' groups related to the effectiveness of EEN.

However, although the sample consisted of severely ill patients and despite the similarities, differences were expected due to patients' illness and to the complexity of the nutritional and to the complexity of the nutritional support. Nutritional support is a complex process involving precise evaluation of the patient's energy requirements, patient's tolerance and the type of underlying disease. Thus, the results showed the following differences: The mean serum albumin value of the burnt patients was significantly increased compared with the multiple trauma patients and the mean serum globulin value of the multiple trauma patients was significantly increased compared with the burnt patients. Furthermore the TIBC increased significantly at the 3rd measurement in burnt patients compared with the multiple trauma patients. The mean value of immunoglobulin IgG showed a significant increase in multiple trauma patients compared with the burnt patients.

Conclusion

Generally, the results of our study confirmed the effectiveness of EEN on both groups of patients as well as the existence of similarities and differences. EEN nutrition provided optimal preservation of the nutritional state of both groups burnt and multiple trauma patients. It is suggested that in severely ill patients nutritional support should be considered as integral part of basic care.

References

1. Wilmore DW, Long JM, Mason AD Jr et al. Catecholamines: mediator of the hypermetabolic response to thermal injury. *Ann Surg* 1974, 180:653-669
2. Bartlett RH, Allyn PA, Medley T et al. Nutritional therapy based on positive caloric balance in burnt patients. *Arch Surg* 1977, 112:974-980
3. Wilmore WD, Mason AD, Johnson W et al. Effect of ambient temperature on heat production and heat loss in burnt patients. *J Appl Physiol* 1975, 38:593-597
4. Alexander JW, Ogle CK, Stinnett JD et al. A sequential, prospective analysis of immunologic abnormalities and infec-

- tion following severe thermal injury. *Ann Surg* 1978, 188:809–816
5. Cheever KH. Early enteral feeding of patients with multiple trauma. *Crit Care Nurse* 1999, 19:40–51
 6. Jeevanandam M, Petersen SR. Substrate fuel kinetics in enterally fed trauma patients supplemented with ornithine alpha ketoglutarate. *Clin Nutr* 1999, 18:209–217
 7. Calder PC. Glutamine and the immune system. *Clin Nutr* 1994, 13:2–8
 8. Wilmore DW, Curreri PW, Spitzer WK et al. Supranormal dietary intake in thermally injured hypermetabolic patients. *Surg Gynecol Obstet* 1971, 132:881–886
 9. Faist E, Schinkel C, Zimmer S. Update on the mechanisms of immune suppression of injury and immune modulation. *World J Surg* 1996, 20:454–459
 10. Jolliet P, Pichard C, Biolo G et al. Enteral nutrition in intensive care patients: A practical approach. Working Group on Nutrition and metabolism, ESICM. European Society of Intensive Care Medicine. *Int Care Med* 1996, 24:848–859
 11. Alexander JW, McMillan BG, Stinnett JD et al. Beneficial effects of aggressive protein feeding in severely burned children. *Ann Surg* 1980, 192:505–507
 12. Jonghe B, appere-De-Vechi C, Fournier M et al. A prospective survey of nutritional support practices in intensive care unit patients: What is prescribed? *Crit Care Med* 2001, 29:8–12
 13. Solem L, Strate RG, Fisher RP. Antacid therapy and nutritional supplementation in the prevention of Curling's ulcer. *Surg Gynecol Obstet* 1979, 148:369–370
 14. Moscona R, Kaufman T, Jacobs R et al. Prevention of gastrointestinal bleeding in burns. The effects of cimetidine or antacids combined with early enteral feeding. *Burns* 1985, 12:65–67
 15. Choctaw WT, Fujita C, Zawacki BE. Prevention of upper gastrointestinal bleeding in burn patients. A role for "elemental" diet. *Arch Surg* 1980, 115:1073–1076
 16. Fadaak HA. Gastrointestinal haemorrhage in burnt patients. The experience of burns unit in Saudi Arabia. *Ann of Burns and Fire Disasters* 2000, XIII:81–83
 17. Raff T, German G, Harman B. The value of early enteral nutrition in the prophylaxis of stress ulceration in the severely burned patient. *Burns* 1997, 23:313–318
 18. Byers PM, Jeejeebhoy KN. Enteral and parenteral nutrition. In: Civetta JM, Talor RW, Kirby RR (eds) *Critical care*. 3rd ed. Philadelphia, Lippincott-Raven Publishers, 1997:464
 19. Adams F, Guest P, Ciraulo D et al. Maximizing tolerance of enteral nutrition in severely injured trauma patients: a comparison of enteral feedings by means of percutaneous endoscopic gastrostomy versus percutaneous endoscopic gastrojejunostomy. *J Trauma* 2001, 48:459–465
 20. Gianotti L, Alexander JW, Nelson JL et al. The role of early enteral feeding and acute starvation on postburn bacterial translocation and host defence: prospective randomized trials. *Crit Care Med* 1994, 22:265–272
 21. Chiarelli A, Enzi G, Casadei A et al. Very early nutrition supplementation in burned patients. *Am J Clin Nutr* 1990, 51:1035–1039
 22. Curreri PW, Richmond D, Marvin et al. Dietary requirements of patients with major burns. *J Am Diet Assoc* 1974, 65:415
 23. Allard JP, Jeejeebhoy KN, Whitwell J et al. Factors influencing energy expenditure in patients with burns. *J Trauma* 1998, 28:199
 24. Deitch EA. The role of intestinal barrier failure and bacterial translocation in the development of systemic infection and multiple organ failure. *Arch Surg* 1990, 125:403–404
 25. Houdijk APJ, Rijnsburger ER, Jansen et al. Randomised trial of glutamine-enriched enteral nutrition on infectious morbidity in patients with multiple trauma. *Lancet* 1998, 352:772–776
 26. Newsholme EA, Crabtree B, Ardawl SM. Glutamine metabolism in lymphocytes: Its biochemical, physiological and clinical importance. *Q J Exp Physiol* 1985, 70:473–499
 27. Liljedahl SO. Treatment of the hypercatabolic state in burns. *Ann Gynaecol* 1980, 69:191–196
 28. Jordan PH, Boulatendis D, Guinn GA. Factors other than vascular occlusion than contribute to intestinal infarction. *Ann Surg* 1970, 171:189–194
 29. Chen LW, Hsu CM, Huang JK et al. Effects of bombesin on gut mucosal immunity in rats after thermal injury. *J Formos Med Assoc* 2000, 6:491–498
 30. Nishimura T, Yamamoto H, DeSerres S et al. Transforming growth factor beta impairs postburn immunoglobulin production by limiting B-cell proliferation, but not cellular synthesis. *J Trauma* 1999, 46:881–885
 31. Enter W, Faist E, Nestle C et al. Dynamics of immunoglobulin synthesis after major trauma. Influence of recombinant lymphokines. *Arch Surg* 1989, 124:1437–1441
 32. Sheridan R, Schulz J, Ryan C et al. Percutaneous endoscopic gastrostomy in burnt patients. *Surg Endosc* 1999, 13:401–402
 33. Curtas S. Closed enteral nutrition delivery systems: fine tuning a safe therapy (Editorial). *Nutrition* 2000, 16:307–308
 34. Kudsk KA, Croce MA, Fabian TC et al. Enteral versus parenteral feeding: effects on septic morbidity after blunt and penetrating abdominal trauma. *Ann Surg* 1992, 215:503–513
 35. Calder PC, Yaqoob P. Glutamine and the immune system. *Amino Acids* 1999, 17:227–241
 36. Cui XL, Iwasa M, Iwasa Y et al. Arginine supplemented diet decreases expression of inflammatory cytokines and improves survival in burned rats. *J Parenter Enteral Nutr* 2000, 24:89–96
 37. Hoodijk APJ, Rijnsburger ER, Jansen J et al. Randomised trial of glutamine-enriched enteral nutrition on infectious morbidity in patients with multiple trauma. *Lancet* 1998, 352:772–776

Αλληλογραφία: Χρ. Μαρβάκη, Αλλαγιάνη 12, 190 03 Μαρκόπουλο Αττικής