Physical activity and energy intake selectively predict the waist-to-hip ratio in men but not in women^{1–3}

Antonia Trichopoulou, Charalambos Gnardellis, Areti Lagiou, Vassiliki Benetou, Androniki Naska, and Dimitrios Trichopoulos

ABSTRACT

Background: The waist-to-hip ratio (WHR) has emerged as an important risk factor for several chronic diseases, but little quantitative information exists about its relation with energy intake and expenditure in men and women.

Objective: We examined the relative role of energy intake and physical activity as determinants of WHRs in men and women, after adjustment for body mass index (BMI) and other likely confounding factors.

Design: In the context of the European Prospective Investigation into Cancer and Nutrition (EPIC), 16433 women and 11520 men aged 30–82 y, apparently healthy and from all over Greece, were examined between 1994 and 1999. Anthropometric measurements were taken, a validated semiquantitative food-frequency questionnaire was administered, and time-weighted occupational and leisure activities were assessed. The WHR was regressed, separately for men and women, on energy intake and energy expenditure after age and BMI were controlled for.

Results: Results for women and men differed. In women, neither energy intake nor energy expenditure was associated with the WHR in any way other than that mediated through BMI. In contrast, in men, higher energy intakes and higher energy expenditures were associated significantly, and largely independently of BMI, with higher and lower WHRs, respectively.

Conclusions: Because the WHR is an important predictor of several cardiovascular and other chronic diseases, documentation of a strong effect of physical activity on the WHR selectively in men may provide a partial explanation of how the effect of physical activity is mediated and why physical activity is more effective in men than in women in reducing disease risk. *Am J Clin Nutr* 2001;74:574–8.

KEY WORDS Body mass index, sex, energy intake, energy expenditure, physical activity, waist-to-hip ratio, Greece

INTRODUCTION

The distribution of adipose tissue in the body, usually derived from the waist-to-hip ratio (WHR), is an important independent risk factor for cardiovascular and possibly other chronic diseases (1, 2). Body mass index (BMI; in kg/m²) is inherently linked to the WHR, and the relation is different in the 2 sexes because the distribution of adipose tissue differs between women and men. Relatively few studies have examined the determinants of the WHR after controlling for BMI. This is an important requirement because BMI and WHR may be independent risk factors for several chronic diseases. Moreover, among all relevant studies only one evaluated the effects of both diet and physical activity in both sexes (3).

We investigated the role of energy intake and expenditure as determinants of the WHR by sex in a large sample of both men and women from the general population. Our study relied on a validated interviewer-administered dietary questionnaire, on detailed information about occupational and recreational physical activity, and on tables of the amount of energy expended with a wide range of physical activities (4, 5).

SUBJECTS AND METHODS

Interviewers administered a validated semiquantitative foodfrequency questionnaire to 27953 apparently healthy men and women aged 30–82 y. The study subjects were recruited from regions throughout Greece to participate in the Greek component of the European Prospective Investigation into Cancer and Nutrition (EPIC) (6). EPIC is a multicenter prospective cohort study investigating the role of nutrition and other lifestyle and environmental factors on the etiology of cancer and other chronic diseases. Approximately 400000 volunteers from 9 European countries participate in the study.

Each participant signed an informed consent form before enrollment. The study was performed in accordance with a protocol approved by the Ethics Committee of the University of Athens. In addition to information regarding nutritional habits, information on lifestyle factors and medical history was

Received July 24, 2000.

Accepted for publication January 16, 2001.

¹From the Department of Hygiene and Epidemiology, University of Athens Medical School, Athens, Greece, and the Department of Epidemiology, Harvard School of Public Health, Boston.

²The European Prospective Investigation into Cancer and Nutrition (EPIC) is coordinated by the International Agency for Research on Cancer and supported by the Europe Against Cancer Programme of the European Commission. The Greek segment of the EPIC study is also supported by the Greek Ministry of Health.

³Address reprint requests to A Trichopoulou, Department of Hygiene and Epidemiology, University of Athens Medical School, 75 Mikras Asias Street, Goudi GR-115 27 Athens, Greece. E-mail: antonia@nut.uoa.gr.

TABLE 1 Waist-to-hip ratio (WHR), energy intake, and energy expenditure by age and sex ¹											
Age (y)	WHR ²		Energy	Energy expenditure ⁴							
	Men	Women	Men	Women	Men	Wom					
			kJ/d		$(MET \cdot h)/d$						
<40 (<i>n</i> = 2081 M, 2766 F)	0.91 ± 0.07	0.76 ± 0.07	11146 ± 3084	8929 ± 2586	36.4 ± 7.0	36.1 ±					
40–54 (<i>n</i> = 4263 M, 5697 F)	0.95 ± 0.07	0.80 ± 0.07	10544 ± 3033	8477 ± 2372	36.3 ± 7.2	36.3 ±					
55–64 ($n = 2404$ M, 4112 F)	0.97 ± 0.07	0.84 ± 0.08	9594 ± 2812	7594 ± 2243	35.5 ± 7.8	35.5 ±					

 0.87 ± 0.08

 0.82 ± 0.08

 0.97 ± 0.07

 0.95 ± 0.07

 $^{1}\overline{x} \pm SD.$

 \geq 65 (*n* = 2772 M, 3858 F)

Total (*n* = 11520 M, 16433 F)

²Comparisons between sexes for all age groups were highly significant (P < 0.01) after Bonferroni adjustment for multiple comparisons. The difference of the mean WHR between the 2 sexes decreased with age, and this interaction was significant (P < 0.01) after Bonferroni adjustment for multiple comparisons (multiple regression with interaction between age and sex).

 8456 ± 2502

 9954 ± 3038

³The sex-by-age interaction was not significant; differences between men and women were highly significant (P < 0.01) after Bonferroni adjustment. There was a significant decrease with age (P < 0.01).

⁴The sex-by-age interaction was not significant; there were no consistent differences between sexes, but there was a highly significant decrease with age (P < 0.01) after Bonferroni adjustment (multiple regression with interaction between age and sex).

recorded through an interviewer-administered lifestyle questionnaire. Anthropometric measurements of weight, height, sitting height, and waist and hip circumferences were taken while the subjects were lightly clothed and wearing no shoes or restrictive underwear. Waist circumference was measured around the smallest circumference between the lowest rib and iliac crest or midway between the lowest rib and iliac crest for obese subjects with no natural waist. The measurement was taken at the end of the normal respiration while the subject was standing erect with his arms at his side and feet together. Hip circumference was measured horizontally at the level of the greatest lateral extension of the hips. All circumferences were measured by using an inelastic tape without compressing the skin and were recorded to the nearest 0.1 cm. The WHR was calculated from the waist and hip measurements. Blood samples were also collected.

The semiquantitative food-frequency questionnaire was validated during the pilot phase of the study (4, 7). The questions covered the average consumption of ≈ 150 food items and beverages over a period of 1 y. Nutrient intakes were calculated through a food-composition database adapted to accommodate the characteristics of the Greek diet (8).

Occupational and leisure time activities were assessed by the use of a special section of the lifestyle questionnaire, which was also used for the overall evaluation of physical activity level. First, the average time spent per day on job-related work of variable intensity, household activities, walking (including walking to work, during shopping, and during leisure time), cycling (including cycling to work and during leisure time), repairing, and gardening (separately for summer and winter) was calculated. The average time spent per day was also assessed for the following leisure-time sporting activities: volleyball, swimming, basketball, soccer, jogging, gymnastics, climbing, dancing, exercise cycling, tennis, rowing, skiing, water skiing, and windsurfing. In assessing sporting activity, we took into account whether the activity was performed throughout the year or seasonally. Finally, the average time spent daily for sleeping (including afternoon naps) and commuting was assessed separately for weekends and weekdays. For subjects whose recorded total hours per day were < or >24 h, the total hours spent daily on each activity were proportionately increased or decreased so that they summed to 24.

Each activity was assigned a MET value (the ratio of the metabolic rate associated with a given activity to the resting metabolic rate) recorded in published tables (5). The time spent in each of the above activities was multiplied by the MET value of that activity, and all MET h products were summed to give a total MET \cdot h score for the day (5, 9). Thus, by assigning h_i to total hours spent per day for an activity *i* with MET value MET_i, the (MET \cdot h)/d score for a set of k activities performed daily, on average, by an individual is defined as follows:

 6791 ± 2059

 7933 ± 2435

$$(\text{MET} \cdot \mathbf{h})/\text{d score} = \Sigma \text{MET}_i \times h_i \tag{1}$$

where i = 1, 2, ..., k. This score essentially corresponds to the number of kilojoules per kilogram body weight expended by an individual during the day.

To assess whether energy intake and expenditure are associated with WHR, after control for the effect of BMI, we regressed WHR (multiplied by 100) on sex, age in years (categorically), total energy intake (in 2092 kJ/d, which corresponds to 500-kcal/d increments, continuously), energy expenditure [in 5-(MET · h)/d increments, continuously], and BMI (continuously). The SPSS 10.0 statistical package (SPSS Inc, Chicago) was used for the statistical analysis.

Regression models were estimated by using data from all 27953 study participants and after exclusion of 8426 participants who reported following a reduced-energy, reduced-salt, or specific macronutrient-reducing diet (notably, a low-fat diet; all such participants are referred to hereafter as dieters), of however short a duration; after exclusion of 7860 study participants who were labeled as underreporters of energy intake; or after exclusion of 13297 study participants who belonged to either of the 2 preceding categories. Study participants were labeled as underreporters of energy intake if their calculated energy intake was $<1.14 \times$ the basal metabolic rate (10–12). The multiple regression with 2 independent factors (age and sex) and their interaction, which was used in the present study, is a standard procedure for applying 2-factor analysis of variance with interaction.

RESULTS

Mean (±SD) WHRs, daily energy intake (kJ), and daily energy expenditure (MET \cdot h) are shown by age and sex in Table 1.

Women

 36.1 ± 4.4

 36.3 ± 4.6

 35.5 ± 5.0

 32.4 ± 4.4

 35.2 ± 4.9

 30.8 ± 5.7

 34.8 ± 7.3

576

TABLE 2

Multiple linear regression-derived coefficients (β), SEs, and *P* values for waist-to-hip ratio (×100) as dependent variable for all participants and after exclusion of those on a diet (A), those who underreported energy intake (B), or A and B^{*I*}

	Men				Women	
Predictor variables	β	SE	Р	β	SE	Р
All participants						
Energy intake (per 2092 kJ/d, 500 kcal/d)	0.16	0.04	< 0.001	-0.07	0.05	0.15
Energy expenditure [per 5 (MET · h)/d]	-0.23	0.04	< 0.001	-0.02	0.06	0.76
BMI (kg/m ²)	0.57	0.01	< 0.001	0.40	0.01	< 0.001
All participants except A						
Energy intake (per 2092 kJ/d, 500 kcal/d)	0.19	0.05	< 0.001	-0.01	0.06	0.83
Energy expenditure [per 5 (MET · h)/d]	-0.24	0.05	< 0.001	0.05	0.07	0.52
BMI (kg/m ²)	0.52	0.01	< 0.001	0.41	0.01	< 0.001
All participants except B						
Energy intake (per 2092 kJ/d, 500 kcal/d)	0.19	0.06	< 0.01	-0.09	0.07	0.16
Energy expenditure [per 5 (MET · h)/d]	-0.29	0.05	< 0.001	-0.02	0.07	0.79
BMI (kg/m ²)	0.59	0.02	< 0.001	0.45	0.01	< 0.001
All participants except A and B ²						
Energy intake (per 2092 kJ/d, 500 kcal/d)	0.22	0.07	< 0.01	-0.07	0.08	0.37
Energy expenditure [per 5 (MET · h)/d]	-0.31	0.06	< 0.001	0.01	0.08	0.95
BMI (kg/m ²)	0.55	0.02	< 0.001	0.45	0.02	< 0.001

¹ Values controlled for age.

²All *P* values for interactions by sex (*F* test) were < 0.01.

WHRs and energy intakes were consistently and significantly higher in men than in women. Energy intake and expenditure decreased significantly with increasing age in both men and women. In contrast, WHRs increased with increasing age in both men and women, but this result was more evident in women (P < 0.01 in all instances). No consistent or significant differences in energy expenditure were noted between the 2 sexes.

Multiple linear regression-derived results for all men and women are shown in Table 2. The results for women and men differed significantly. In women, neither energy intake nor energy expenditure was associated with WHR in any way other than that mediated through BMI. BMI was an important predictor of WHR; for example, in women who were neither dieters nor underreporters of energy intake, a BMI that was 1 kg/m² higher corresponded to a WHR (multiplied by 100) that was higher by a value of 0.45 (95% CI: 0.42, 0.48). For men, the results were strikingly different. BMI was associated with WHR more strongly in men than in women. For example, in men who were neither dieters nor underreporters of energy intake, the regression coefficient was 0.55 (95% CI: 0.52, 0.58). The difference between 0.45 and 0.55 was significant (*P* for interaction < 0.001). More importantly, and in contrast with the finding for women, in men, higher energy intakes and higher energy expenditures were associated significantly, and largely independently of BMI, with higher and lower WHRs, respectively.

We also evaluated the overall effects of energy intake and energy expenditure on WHR, whether these effects were independent of BMI (as in Table 2) or were mediated through BMI. After deletion of the BMI variable from the models that generated the results shown in Table 2 and after exclusion of both underreporters and dieters, there was a weak and nonsignificant negative effect of energy expenditure in women, whereas energy intake was significantly positively associated with WHR (data not shown). In contrast, and as expected, in men the regression coefficient for energy intake increased from 0.22 to 0.57, whereas the regression coefficient for energy expenditure decreased from -0.31 to -0.37. Finally, when we ran the models in Table 2 with the inclusion of tobacco smoking as an additional predictor variable, the regression coefficients remained essentially unchanged. This finding indicates that tobacco smoking was not an important confounder of the studied correlations, despite its known relation with body weight.

DISCUSSION

The results of this large investigation indicated that energy intake and expenditure had essentially no association with WHR in women when BMI was controlled for. In other words, in women, higher energy intakes and higher energy expenditures corresponded to slightly higher and slightly lower WHRs, respectively, but both these effects were largely mediated through BMI, which is known to reflect the balance of energy intake and expenditure (13). On the contrary, in men, higher energy intakes and higher energy expenditures corresponded to significantly higher and significantly lower WHRs, respectively, even when BMI was accounted for. In addition, whereas reductions in BMI are more efficiently achieved with reductions in energy intake than with increases in energy expenditure in both sexes, the opposite appeared to be true for a reduction of WHR in men (13).

This investigation had the weaknesses inherent in crosssectional investigations that have to account for energy intake and expenditure as well as BMI (13, 14). Our study, however, also had considerable advantages: the study was large, 5 times as large as any other investigation of the differential role of energy intake and expenditure on WHR by sex (3); the dietary questionnaire was validated and administered by specially trained interviewers (4, 7); and the assessment of physical activity relied on many questions covering the type, intensity, and duration of a broad range of activities. The physical activity questionnaire was also validated originally, albeit in a different population (15). From another point of view, each of our principal results was compatible with the findings of previous investigations (16–30), a fact that supports the validity of our results. Our study, however, was both larger and of broader scope than most previous investigations that focused on either energy intake or energy expenditure or physical activity.

Ever since WHR was proposed as an important independent risk factor for many chronic diseases (1, 31), several studies have indicated that changes in waist circumference and hip circumference and the corresponding ratio are associated with conditions such as type 2 diabetes (32-35), hypertension and stroke (33, 36), dislipidemias (33, 37), and benign prostatic hypertrophy (2). Although all of the studies did not confirm all aspects of the preceding constellation of findings (38), the WHR is now generally recognized as an important risk factor, possibly linked with the phenomenon of insulin resistance and the insulin-like-growthfactor system. Several studies evaluated the effect of energy expenditure on WHR in women (16, 17), men (18-21), or both sexes (22-27), whereas others have evaluated the effects of both energy intake and expenditure on WHR in women (28) or men (29, 30). To our knowledge, however, only the study by Slattery et al (3) evaluated the effects of both diet and physical activity in both men and women. In comparison with that pioneer study, our investigation was 5 times larger and relied on an energy expenditure metric $[(MET \cdot h)/d]$ that allows comparisons to be made between energy intake and expenditure.

Although the information in the existing literature on the relation between WHR and energy intake and expenditure is fragmentary, was obtained with different methods, and relied on different population samples, it is fairly consistent in indicating that, after BMI is controlled for, increases in energy intake and decreases in energy expenditure sharply reduce WHRs in men but not in women (25). These findings are intuitively appealing because physically inactive, middle-aged men are frequently recognized by their protruding abdomen. The findings, however, may have important implications, including an explanation of why physical activity may be more important for health in men than in women.

REFERENCES

- Daly PA, Solomon CG, Manson J. Risk modification in the obese patient. In: Manson J, Ridker P, Gaziano JM, Hennekens C, eds. Prevention of myocardial infarction. New York: Oxford University Press, 1996:231–63.
- Giovannucci E, Rimm ER, Chute CG, et al. Obesity and benign prostatic hyperplasia. Am J Epidemiol 1994;140:989–1002.
- Slattery ML, McDonald A, Bild DE, et al. Associations of body fat and its distribution with dietary intake, physical activity, alcohol, and smoking in blacks and whites. Am J Clin Nutr 1992;55:943–9.
- Gnardellis C, Trichopoulou A, Katsouyianni K, Polychronopoulos E, Rimm EB, Trichopoulos D. Reproducibility and validity of an extensive semiquantitative food frequency questionnaire among Greek school teachers. Epidemiology 1995;6:74–7.
- Ainsworth BE, Haskell WL, Leon AS. Compendium of physical activities: classification of energy costs of human physical activities. Med Sci Sports Exerc 1993;25:71–80.
- Riboli E. Nutrition and cancer: background and rationale of the European Prospective Investigation into Cancer and Nutrition (EPIC). Ann Oncol 1992;3:783–91.
- Katsouyanni K, Rimm EB, Gnardellis C, Trichopoulos D, Polychronopoulos E, Trichopoulou A. Reproducibility and relative validity of an extensive semi-quantitative food frequency questionnaire using dietary records and biochemical markers among Greek schoolteachers. Int J Epidemiol 1997;26(suppl):S118–27.
- Trichopoulou A. Composition of Greek foods and dishes. Athens, Greece: National School of Public Health, 1992.

- Martinez ME, Giovannucci E, Spiegelman D, Hunter DJ, Willett WC, Colditz GA. Leisure-time physical activity, body size, and colon cancer in women. J Natl Cancer Inst 1997;89:948–55.
- Goldberg GR, Black AE, Jebb SA, et al. Critical evaluation of energy intake data using fundamental principles of energy physiology: 1. Derivation of cut-off values to identify under-recording. Eur J Clin Nutr 1991;45:569–81.
- Gnardellis C, Boulou C, Trichopoulou A. Magnitude determinants and impact of under-reporting of energy intake in a cohort study in Greece. Public Health Nutr 1998;1:131–7.
- Shetty PS, Henry CJ, Black AE, Prentice AM. Energy requirements of adults: an update on basal metabolic rates (BMRs) and physical activity levels (PALs). Eur J Clin Nutr 1996;50(suppl):S11–23.
- Trichopoulou A, Gnardellis C, Lagiou A, Benetou V, Trichopoulos D. Body mass index in relation to energy intake and expenditure among adults in Greece. Epidemiology 2000;11:333–6.
- Willett WC. Nutritional epidemiology. New York: Oxford University Press, 1998.
- Pols MA, Peeters PHM, Cocke M, Slimani N, Bueno-de-Mesquita HB, Collette HJA. Estimation of reproducibility and relative validity of the questions included in the EPIC physical activity questionnaire. Int J Epidemiol 1997;26(suppl):S181–9.
- Pols MA, Peeters PH, Twisk JW, Kemper HC, Grobbee DE. Physical activity and cardiovascular disease risk profile in women. Am J Epidemiol 1997;146:322–8.
- Davy KP, Evans SL, Stevenson ET, Seals DR. Adiposity and regional body fat distribution in physically active young and middle-aged women. Int J Obes Relat Metab Disord 1996;20:777–83.
- Selby JV, Newman B, Quesenberry CP, et al. Genetic and behavioral influences on body fat distribution. Int J Obes 1990;14:593–602.
- Seidell JC, Cibolini M, Deslypere JP, Charzewska J, Ellsinger BM, Cruz A. Body fat distribution in relation to physical activity and smoking habits in 38-year-old European men. The European Fat Distribution Study. Am J Epidemiol 1991;133:257–65.
- Houmard JA, McCulley C, Roy LK, Bruner RK, McCammon MR, Israel RG. Effects of exercise training on absolute and regional adiposity. Int J Obes Relat Metab Disord 1994;18:243–8.
- Nylind B, Schele R, Linroth K. Changes in male exercise performance and anthropometric variables between the ages of 19 and 30. Eur J Appl Physiol 1978;38:145–50.
- 22. Wilmore JH, Despres JP, Stanforth PR, et al. Alterations in body weight and composition consequent to 20 wk of endurance training: the HERITAGE Family Study. Am J Clin Nutr 1999;70:346–52.
- 23. Kriska AM, LaPorte RE, Pettitt DJ, et al. The association of physical activity with obesity, fat distribution and glucose intolerance in Pima Indians. Diabetologia 1993;36:863–9.
- Visser M, Launer LJ, Deurenberg P, Deeg DJ. Total and sports activity in older men and women: relation with body fat distribution. Am J Epidemiol 1997;145:752–61.
- Ross R. Effects of diet- and exercise-induced weight loss on visceral adipose tissue in men and women. Sports Med 1997;24:55–64.
- Ross R, Janssen I. Is abdominal fat preferentially reduced in response to exercise-induced weight loss? Med Sci Sports Exerc 1999;31(suppl):S568–72.
- Lehmann R, Vokac A, Niedermann K, Agosti K, Spinas GA. Loss of abdominal fat and improvement of the cardiovascular risk profile by regular moderate exercise training in patients with NIDDM. Diabetologia 1995;38:1313–9.
- Rose KM, Newman B, Mayer Davis EJ, Selby JV. Genetic and behavioral determinants of waist-hip ratio and waist circumference in women twins. Obes Res 1998;6:383–92.
- Troisi RJ, Heinold JW, Vokonas PS, Weiss ST. Cigarette smoking, dietary intake, and physical activity: effects on body fat distribution—the Normative Aging Study. Am J Clin Nutr 1991;53:1104–11.
- 30. Hellinius ML, De Fair U, Berlung B, Hamsten A, Krakau I. Diet and exercise are equally effective in reducing risk for cardiovascular disease. Results of a randomized control study in men with slightly to

moderately raised cardiovascular risk factors. Atherosclerosis 1993;103:81-91.

- 31. Lapidus L, Bengtsson C, Larsson B, et al. Distribution of adipose tissue and risk of cardiovascular disease and death: a 12 year follow up of participants in the population study of women in Gothenberg, Sweden. Br Med J (Clin Res Ed) 1984;289:1257–61.
- 32. Carey VJ, Walters EE, Colditz GA, et al. Body fat distribution and risk of non-insulin-dependent diabetes mellitus in women. The Nurses' Health Study. Am J Epidemiol 1997;145:614–9.
- Ledoux M, Lambert J, Reeder BA, Despres JP. Correlation between cardiovascular disease risk factors and simple anthropometric measures. Canadian Heart Health Surveys Research Group. CMAJ 1997;157(suppl):S46–53.
- 34. Seidell JC, Han TS, Feskens EJ, Lean ME. Narrow hips and broad waist circumferences independently contribute to increased risk of

non-insulin-dependent diabetes mellitus. J Intern Med 1997;242: 401-6.

- 35. Okosun IS, Cooper RS, Rotimi CN, Osotimehin B, Forrester T. Association of waist circumference with risk of hypertension and type 2 diabetes in Nigerians, Jamaicans, and African-Americans. Diabetes Care 1998;21:1836–42.
- Folsom AR, Prineas RJ, Kaye SA, Munger RG. Incidence of hypertension and stroke in relation to body fat distribution and other risk factors in older women. Stroke 1990;21:701–6.
- Lehmann R, Kaplan V, Bingisser R, Bloch KE, Spinas GA. Impact of physical activity on cardiovascular risk factors in IDDM. Diabetes Care 1997;20:1603–11.
- Bonora E, Zenere M, Branzi P, et al. Influence of body fat and its regional localization on risk factors for atherosclerosis in young men. Am J Epidemiol 1992;135:1271–8.